

Package: DiscreteLaplace (via r-universe)

December 25, 2024

Type Package

Title Discrete Laplace Distributions

Version 1.1.1

Date 2016-04-29

Author Alessandro Barbiero <alessandro.barbiero@unimi.it>, Riccardo Inchingolo <dott.inchingolo_r@libero.it>

Maintainer Alessandro Barbiero <alessandro.barbiero@unimi.it>

Description Probability mass function, distribution function, quantile function, random generation and estimation for the skew discrete Laplace distributions.

License GPL

LazyLoad yes

Date/Publication 2016-05-01 00:46:21

NeedsCompilation no

Repository <https://alessandro-barbiero.r-universe.dev>

RemoteUrl <https://github.com/cran/DiscreteLaplace>

RemoteRef HEAD

RemoteSha 898b019fd0b368e8268dd8f1433a3b57e3f6a29e

Contents

DiscreteLaplace-package	2
ddlaplace	3
ddlaplace2	4
dlaplacelike2	6
Edlaplace	7
Edlaplace2	8
estdlaplace	9
estdlaplace2	11
iFI	12
iFI2	13
ioFI2	14
loss	15

DiscreteLaplace-package
Discrete Laplace Distributions

Description

Probability mass function, distribution function, quantile function, random generation and sample estimation for two discrete skew Laplace distributions on integers. The skew discrete Laplace distributions here considered are that proposed by Kozubowski and Inusah (2006), henceforth referred to as DSL, and the alternative one proposed by Barbiero (2014), henceforth ADSL.

Details

Package:	DiscreteLaplace
Type:	Package
Version:	1.1.1
Date:	2016-04-29
License:	GPL
LazyLoad:	yes

Author(s)

Alessandro Barbiero, Riccardo Inchingolo

Maintainer: Alessandro Barbiero <alessandro.barbiero@unimi.it>

References

T. J. Kozubowski, S. Inusah (2006) A skew Laplace distribution on integers, *Annals of the Institute of Statistical Mathematics*, 58: 555-571, <http://dx.doi.org/10.1007/s10463-005-0029-1>

A. Barbiero (2014) An alternative discrete Laplace distribution, *Statistical Methodology*, 16: 47-67, <http://dx.doi.org/10.1016/j.stamet.2013.07.002>

ddlaplace	<i>Probability mass function, cumulative distribution function, quantile function and random generation of the DSL</i>
-----------	--

Description

The function computes the probability mass function, the cumulative distribution function, the quantile function of the DSL and implements random generation.

Usage

```
ddlaplace(x, p, q)
pdlaplace(x, p, q)
qdlaplace(prob, p, q)
rdlaplace(n, p, q)
```

Arguments

x	vector of quantiles
p	the first parameter p in $(0, 1)$ of the DSL
q	the second parameter q in $(0, 1)$ of the DSL
prob	vector of probabilities
n	number of observations

Details

The pmf of the DSL is given by

$$P(X = x; p, q) = \frac{(1-p)(1-q)}{1-pq} p^x; x = 0, 1, 2, 3, \dots$$

$$P(X = x; p, q) = \frac{(1-p)(1-q)}{1-pq} q^{|x|}; x = 0, -1, -2, -3, \dots$$

whereas the cumulative distribution function is given by

$$F(x; p, q) = P(X \leq x) = \frac{(1-p)q^{-\lfloor x \rfloor}}{1-pq}, x < 0$$

$$F(x; p, q) = P(X \leq x) = 1 - \frac{(1-q)p^{\lfloor x \rfloor + 1}}{1-pq}, x \geq 0$$

Value

ddlaplace returns the probability of x; pdlaplace returns the cumulate probability of x; qdlaplace returns the prob- quantile; rdlaplace returns a random sample of size n from DSL.

Author(s)

Alessandro Barbiero, Riccardo Inchingolo

References

Tomasz J. Kozubowski, Seidu Inusah (2006) A skew Laplace distribution on integers, *Annals of the Institute of Statistical Mathematics*, 58: 555-571

See Also

[ddlaplace2](#)

Examples

```
# pmf
p<-0.7
q<-0.45
x<-10:10
prob<-ddlaplace(x, p, q)
plot(x, prob, type="h")
prob<-ddlaplace(x, q, p) # swap the parameters
plot(x, prob, type="h")
ddlaplace(-4:4, 1:9/10, 9:1/10) # letting p and q be vectors
# cdf
p<-0.2
q<-0.5
x<-c(-3, -1, pi)
pdlaplace(x, p, q)
# quantile function
p<-0.8
q<-0.4
prob<-c(0.2,0.5,0.8)
x<-qdlaplace(prob, p, q)
x # check
upper<-pdlaplace(x, p, q)
upper
lower<-pdlaplace(x-1, p, q)
lower
lower<=prob & prob<=upper
# random generation
n<-100
p<-0.3
q<-0.5
x<-rdlaplace(n, p, q)
x
t<-table(x)
t
plot(t)
```

Description

The function computes the probability mass function, the cumulative distribution function, the quantile function of the ADSL and provides random generation of samples from the same model

Usage

```
ddlaplace2(x, p, q)
palaplace2(x, p, q)
pdlaplace2(x, p, q)
qdlaplace2(prob, p, q)
rdlaplace2(n, p, q)
```

Arguments

x	vector of quantiles
p	the first parameter p in $(0, 1)$ of the ADSL
q	the second parameter q in $(0, 1)$ of the ADSL
prob	vector of probabilities
n	number of observations

Details

The probability mass function of the ADSL distribution is given by:

$$P(X = x; p, q) = \frac{\log p}{\log(pq)} q^{-(x+1)}(1-q) \text{ for } x = \dots, -2, -1$$

and

$$P(X = x; p, q) = \frac{\log q}{\log(pq)} p^x(1-p) \text{ for } x = 0, 1, 2, \dots$$

Its cumulative distribution function is:

$$F(x; p, q) = \frac{\log p}{\log(pq)} q^{-(\lfloor x \rfloor + 1)} \text{ for } x < 0$$

and

$$F(x; p, q) = 1 - \frac{\log q}{\log(pq)} p^{(\lfloor x \rfloor + 1)} \text{ for } x \geq 0$$

Value

ddlaplace2 returns the probability of x; pdlaplace2 returns the cumulative probability of x; qdlaplace2 returns the prob- quantile; rdlaplace2 returns a random sample of size n from ADSL.

Author(s)

Alessandro Barbiero, Riccardo Inchingolo

References

A. Barbiero, An alternative discrete Laplace distribution, *Statistical Methodology*, 16: 47-67

See Also[ddlaplace](#)**Examples**

```
# pmf
p <- 0.7
q <- 0.45
x <- -10:10
prob <- dlaplace2(x, p, q)
plot(x, prob, type="h")
# swap the parameters
prob <- dlaplace2(x, q, p)
plot(x, prob, type="h")
# letting p and q be vectors...
ddlaplace2(-4:4, 1:9/10, 9:1/10)
# cdf
pdlaplace2(x, p, q)
pdlaplace2(pi, p, q)
pdlaplace2(floor(pi), p, q)
# quantile function
qdlaplace(1:9/10, p, q)
# random generation
y <- rdlaplace2(n=1000, p, q)
plot(table(y))
```

dlaplace2*Log-likelihood function for the ADSL distribution***Description**

Log-likelihood function (changed in sign) for the ADSL distribution.

Usage

```
dlaplace2(par, x)
```

Arguments

<code>par</code>	the vector of parameters (p, q)
<code>x</code>	a vector of observations from ADSL

Value

The log-likelihood function with changed sign.

Author(s)

Alessandro Barbiero, Riccardo Inchingolo

References

A. Barbiero, An alternative discrete Laplace distribution, *Statistical Methodology*, 16: 47-67

See Also

[estdlaplace2](#), [ddlaplace2](#)

Examples

```
p <- 0.25
q <- 0.7
x <- rdlaplace2(n=100, p, q)
par <- estdlaplace2(x, "ML")
-dlaplacelike2(par, x) # greater than...
-dlaplacelike2(c(p, q), x)
```

Edlaplace

Moments of the discrete Laplace distribution

Description

The function provides the expected value and the variance of the SDL, and the expectation of its absolute value.

Usage

`Edlaplace(p, q)`

Arguments

- p the first parameter, in (0, 1), of the DSL
- q the second parameter, in (0, 1), of the DSL

Details

$$\begin{aligned} E(X; p, q) &= \frac{1}{1-p} - \frac{1}{1-q} = \frac{p}{1-p} - \frac{q}{1-q}, \\ E(|X|; p, q) &= \frac{q(1-p)^2 + p(1-q)^2}{(1-qp)(1-q)(1-p)}, \\ V(X; p, q) &= \frac{1}{(1-p)^2(1-q)^2} \left[\frac{q(1-p)^3(1+q) + p(1-q)^3(1+p)}{1-pq} - (p-q)^2 \right] \end{aligned}$$

Value

A list of three items:

- E1 expected value
- E1a expectation of the absolute value
- V variance

Author(s)

Alessandro Barbiero, Riccardo Inchingolo

References

T. J. Kozubowski, S. Inusah (2006) A skew Laplace distribution on integers, *Annals of the Institute of Statistical Mathematics*, 58: 555-571

See Also

[ddlaplace](#)

Examples

```
# ex.1
p<-0.5
q<-0.4
Edlaplace(p, q)
# ex.2
p<-0.1
q<-0.9
Edlaplace(p, q)
```

Edlaplace2

First- and second-order moments of ADSL

Description

First- and second-order moment of the ADSL distribution.

Usage

Edlaplace2(p, q)

Arguments

- | | |
|---|---|
| p | the first parameter p , in $(0, 1)$, of the ADSL |
| q | the first parameter q , in $(0, 1)$, of the ADSL |

Details

For the ADSL distribution,

$$E(X; p, q) = \frac{\log q}{\log(pq)} \frac{p}{1-p} - \frac{\log p}{\log(pq)} \frac{1}{1-q}$$

and

$$E(X^2; p, q) = \frac{\log q}{\log(pq)} \frac{p(1+p)}{(1-p)^2} + \frac{\log p}{\log(pq)} \frac{1+q}{(1-q)^2}$$

Value

A list containing the first- and the second-order moments of the ADSL distribution, E1 and E2.

Author(s)

Alessandro Barbiero, Riccardo Inchingolo

References

A. Barbiero, An alternative discrete Laplace distribution, *Statistical Methodology*, 16: 47-67

See Also

[estdlaplace2](#), [loss](#), [ddlaplace2](#)

Examples

```
Edlaplace2(p=0.3, q=0.3)
Edlaplace2(p=0.3, q=0.6)
Edlaplace2(p=0.6, q=0.3)
Edlaplace2(p=0.6, q=0.6)
```

estdlaplace

Sample estimation for the DSL

Description

The function provides the maximum likelihood estimates for the parameters of the DSL and the estimate of the inverse of the Fisher information matrix. The method of moments estimates of p and q coincide with the maximum likelihood estimates.

Usage

`estdlaplace(x)`

Arguments

`x` a vector of observations from the DSL

Details

See the reference. If $\bar{x}^+ = \frac{1}{n} \sum_{i=1}^n x_i^+$, $\bar{x}^- = \frac{1}{n} \sum_{i=1}^n x_i^-$ where x^+ and x^- are the positive and the negative parts of x , respectively: $x^+ = x$ if $x \geq 0$ and zero otherwise, $x^- = (-x)^+$, then

$$\hat{q} = \frac{2\bar{x}^-(1+\bar{x})}{1+2\bar{x}^-\bar{x}+\sqrt{1+4\bar{x}^-\bar{x}^+}}, \hat{p} = \frac{\hat{q}+\bar{x}(1-\hat{q})}{1+\bar{x}(1-\hat{q})}$$

when $\bar{x} \geq 0$ and

$$\hat{p} = \frac{2\bar{x}^+(1-\bar{x})}{1-2\bar{x}^+\bar{x}+\sqrt{1+4\bar{x}^-\bar{x}^+}}, \hat{q} = \frac{\hat{p}-\bar{x}(1-\hat{p})}{1-\bar{x}(1-\hat{p})}$$

when $\bar{x} \leq 0$.

Value

A list comprising

<code>hatp</code>	estimate of p
<code>hatq</code>	estimate of q
<code>hatSigma</code>	estimate of the inverse of the Fisher information matrix

Author(s)

Alessandro Barbiero, Riccardo Inchingolo

References

T. J. Kozubowski, S. Inusah (2006) A skew Laplace distribution on integers, *Annals of the Institute of Statistical Mathematics*, 58: 555-571

See Also

[ddlaplace](#)

Examples

```
p<-0.6
q<-0.3
n<-20
x<-rdlaplace(n, p, q)
est<-estdlaplace(x)
est[1]
est[2]
est[3]
# increase n
n<-100
x<-rdlaplace(n, p, q)
est<-estdlaplace(x)
est[1]
est[2]
est[3]
# swap the parameters
x<-rdlaplace(n, q, p)
est<-estdlaplace(x)
est[1]
est[2]
est[3]
```

estdlaplace2*Sample estimation for the ADSL***Description**

The function provides the point estimates for the parameters of the ADSL, resorting to four possible methods: method of moments, maximum likelihood method, method of proportion, modified method of moments. For details, please take a look at the references.

Usage

```
estdlaplace2(x, method = "M", err = 0.001, parml = c(exp(-1), exp(-1)))
```

Arguments

x	a vector of observations from the ADSL
method	M for the method of moments, ML for the maximum likelihood methods, P for the method of proportion, MM for the modified method of moments
err	a positive tolerance value, as small as possible, used in the definition of lower and upper bounds of the parameters p and q in the minimization algorithm utilized by the method of moments
parml	starting values for p and q in the optimization process for the maximum likelihood method

Value

a vector with the parameter estimates of p and q .

Author(s)

Alessandro Barbiero, Riccardo Inchingolo

References

A. Barbiero, An alternative discrete Laplace distribution, *Statistical Methodology*, 16: 47-67

See Also

[dlaplace2](#)

Examples

```
p <- 0.4
q <- 0.6
x <- rdlaplace2(n=100, p, q)
est <- matrix(0, 5, 2)
est[1,] <- c(p,q)
est[2,] <- estdlaplace2(x, method="M")
```

```

est[3,] <- estdlaplace2(x, method="ML")
est[4,] <- estdlaplace2(x, method="P")
est[5,] <- estdlaplace2(x, method="MM")
dimnames(est)[[1]]<-c("true","M","ML","P","MM")
dimnames(est)[[2]]<-c("p","q")
xlim <- c(min(est[,1])* .98,max(est[,1])*1.02)
ylim <- c(min(est[,2])* .98,max(est[,2])*1.02)
plot(est, pch=19, col=1:5, xlim=xlim, ylim=ylim)
text(est, dimnames(est)[[1]], pos=3, col=1:5, cex= .75)

```

iFI*Inverse of Fisher Information matrix***Description**

Inverse of Fisher Information matrix for the DSL.

Usage

```
iFI(p, q)
```

Arguments

- | | |
|----------|---|
| p | first parameter p , in $(0, 1)$, of the DSL |
| q | second parameter q , in $(0, 1)$, of the DSL |

Value

The inverse of Fisher Information matrix. Take a look at the references for more details.

Author(s)

Alessandro Barbiero, Riccardo Inchingolo

References

- T. J. Kozubowski, S. Inusah (2006) A skew Laplace distribution on integers, *Annals of the Institute of Statistical Mathematics*, 58: 555-571

See Also

[iFI2](#)

Examples

```

p <- 0.2
q <- 0.8
iFI(p, q)

```

iFI2

Inverse of Fisher Information matrix

Description

Inverse of Fisher Information matrix for the ADSL.

Usage

`iFI2(p, q)`

Arguments

- | | |
|----------------|--|
| <code>p</code> | first parameter p , in $(0, 1)$, of the ADSL |
| <code>q</code> | second parameter q , in $(0, 1)$, of the ADSL |

Value

The inverse of Fisher Information matrix. Take a look at the references for more details.

Author(s)

Alessandro Barbiero, Riccardo Inchingolo

References

A. Barbiero, An alternative discrete Laplace distribution, *Statistical Methodology*, 16: 47-67

See Also

[ioFI2](#)

Examples

```
p <- 0.2  
q <- 0.8  
iFI2(p, q)
```

ioFI2*Inverse of the observed Fisher Information matrix***Description**

Inverse of the observed Fisher Information matrix computed on a random sample of ADSL values.

Usage

```
ioFI2(x)
```

Arguments

<code>x</code>	a vector of observations from the ADSL
----------------	--

Value

The inverse of the observed Fisher Information matrix.

Author(s)

Alessandro Barbiero, Riccardo Inchingolo

References

A. Barbiero, An alternative discrete Laplace distribution, *Statistical Methodology*, 16: 47-67

See Also

[iFI2](#)

Examples

```
n <- 100
p <- 0.4
q <- 0.7
x <- rdlaplace2(n, p, q)
M <- ioFI2(x)
par <- estdlaplace2(x, "ML")
se <- diag(sqrt(M))
par # MLEs
se # standard errors
M # compare with the inverse of Fisher Information matrix
iFI2(par[1], par[2])/n # with MLEs plugged in
iFI2(p, q)/n # or the true values
```

loss	<i>Loss function for the method of moments</i>
------	--

Description

A loss function used for the implementation of the method of moments (for the ADSL).

Usage

```
loss(par, x)
```

Arguments

- | | |
|-----|--|
| par | the vector of parameters, p and q, of the ADSL |
| x | a vector of sample values from the ADSL |

Value

The value $L = [E(X) - m_1(x)]^2 + [E(X^2) - m_2(x)]^2$, where m_1 and m_2 are the first- and second-order sample moments.

Author(s)

Alessandro Barbiero, Riccardo Inchingolo

References

A. Barbiero, An alternative discrete Laplace distribution, *Statistical Methodology*, 16: 47-67

See Also

[estdlaplace2](#), [Edlaplace2](#)

Examples

```
p <- 0.3
q <- 0.7
x <- rdlaplace2(n=100, p, q)
par <- estdlaplace2(x, "M")
loss(par, x) # should be near zero
loss(c(p,q), x) # may be far greater than zero
```

Index

* **distribution,htest**
 estdlaplace2, 11
 iFI, 12
 iFI2, 13
* **distribution**
 ddlaplace, 3
 ddlaplace2, 4
 Edlaplace, 7
 Edlaplace2, 8
 estdlaplace, 9
 ioFI2, 14
 loss, 15
* **htest**
 dlaplace2, 6
 estdlaplace, 9
 ioFI2, 14
* **package**
 DiscreteLaplace-package, 2

ddlaplace, 3, 6, 8, 10
ddlaplace2, 4, 4, 7, 9
DiscreteLaplace
 (DiscreteLaplace-package), 2
DiscreteLaplace-package, 2
dlaplace2, 6, 11

 Edlaplace, 7
 Edlaplace2, 8, 15
 estdlaplace, 9
 estdlaplace2, 7, 9, 11, 15

 iFI, 12
 iFI2, 12, 13, 14
 ioFI2, 13, 14

 loss, 9, 15

palaplace2 (ddlaplace2), 4
pdlaplace (ddlaplace), 3
pdlaplace2 (ddlaplace2), 4